

## CORNEAL ULCERS

The cornea is the clear, shiny membrane which makes up the surface of the eyeball. It is much like a clear window. To understand a corneal ulcer, you must first understand how the cornea is constructed.

The cornea is comprised of three layers. The most superficial layer, the epithelium, is comprised of many very thin layers of cells. Below the epithelium is the stroma, and the deepest layer is Descemet's membrane. Because all of these layers are clear, it is not possible to see them without special stains and a microscope.

A shallow scrape through a few layers of the epithelium is called a *corneal erosion* or a *corneal abrasion*. A *corneal ulcer* is erosion through the entire epithelium and into the stroma. If the erosion goes through the epithelium and stroma to the level of Descemet's membrane, a *descemetocoele* exists. If Descemet's membrane ruptures, the liquid inside the eyeball leaks out and the eye collapses.

### ***What causes corneal ulcers?***

There are several causes for corneal ulcers in cats. The most common is trauma. An ulcer may result from blunt trauma, such as a cat rubbing its eye on a carpet, or due to a laceration, such as a cat scratch.

Other causes of corneal ulcers include bacterial infections, viral infections, and other diseases. These may originate in the eye or develop secondary to disease elsewhere in the body.

### ***What will my cat do if an ulcer is present?***

A corneal ulcer is very painful. In response to pain, most cats rub the affected eye with a foot or on the carpet. To protect the eye, they keep the lids tightly closed. Occasionally, there will be a discharge that collects in the corner of the eye or runs down the face.

### ***How is a corneal ulcer diagnosed?***

Superficial corneal abrasions are usually not visible. They can be highlighted and seen with the use of fluorescein stain. A drop of this orange-colored stain is placed on the cornea. The dye will adhere to an area of ulceration and is easily visualized with a special black light called a Wood's light. This is the most basic test performed and may be the only test needed if the ulcer is acute and very superficial. If the ulcerated area is chronic or very deep, samples may be taken for culture and cell study prior to applying the stain or any other medication.

### ***Can an ulcer be treated?***

Yes. However, the form of treatment depends on whether there is a corneal abrasion, corneal ulcer, or descemetocoele present.

Corneal abrasions generally heal within 3-5 days. Medication is used to prevent bacterial infections (antibiotic ophthalmic drops or ointment) and to relieve pain (atropine ophthalmic drops or ointment). Antibiotic drops are only effective for a few minutes so they must be applied frequently; ointments last a bit longer but still require application every few hours. It is suggested that an antibiotic preparation be instilled in the eye four to six times per day. On the other hand, the effects of atropine last many hours so this drug is only used twice daily.

If a corneal ulcer or descemetocoele is present, measures must be taken to protect the eye and to promote healing. Since cats do not wear eye patches well, surgical techniques are often used to close the eyelids

and cover the ulcer or descemetocele. These measures protect the eye for several days, then are reversed so the cat can use the eye again.

Ulcers that do not heal well often have a buildup of dead cells at the ulcer edge. These dead cells prevent normal cells from the corneal surface from sliding over the ulcer edge and filling in the defect. If this appears to be part of the healing problem, the dead cells are removed from the edges of the ulcer before the eyelids are surgically closed. In some cases, removing the dead cells may be all that is needed to start the healing process, so surgical closing of the eyelids may not be necessary.

***Is there a difference between a corneal ulcer and a corneal abrasion?***

It is easy to confuse an ulcer and an abrasion on the first examination. There is a judgment call involved in differentiating the two. After 2-3 days of treatment, your cat should be reexamined to be sure that healing is progressing properly. If that does not happen, a decision may be made to perform surgery.

***Do any of the medications have side-effects?***

Rarely, a cat will be allergic to an antibiotic that is instilled in the eye. If your cat seems to be in more pain after the medication is used, discontinue it and contact the veterinarian.

A cat with a corneal ulcer has quite a bit of pain in the eye, so it keeps it tightly shut. Atropine is used to relieve that pain. However, atropine also dilates the pupil widely. This means that the cat is very sensitive to light in that eye. Because of the light sensitivity, the eye will be held closed in bright light.

Atropine's effects may last for several days after the drug is discontinued. Do not be alarmed if the pupil stays dilated for several days. Should you accidentally get atropine in your eye, the same prolonged pupillary dilation will occur.

Atropine has a very objectionable taste. If it gets in the cat's mouth, drooling will occur. This is not a drug reaction but a reaction to the terrible taste. It will subside in a few minutes. However, once this happens, the cat may be more difficult to medicate because it may anticipate the same thing happening again.

***What can I use to control the pain?***

A topical anesthetic is often used to numb the cornea so diagnostic tests may be performed. However, these drugs are toxic to the corneal epithelium and prevent proper healing. They are safe for one time use, but they should not be used as part of the treatment.

***How do I know when the treatment is completed?***

The best way to tell that the cornea has healed is to repeat the fluorescein stain test. This should be done after approximately 5-7 days of treatment.

***What does it mean if there are red streaks near the ulcer?***

The normal cornea has no blood vessels going through it. However, when a corneal ulcer or descemetocele occurs, the body senses a need to increase its healing capabilities. New blood vessels are created by a process called neovascularization. The new vessels begin at the sclera (the white part of the eye) and course their way to the ulcer.